European Journal of Pharmacology – Molecular Pharmacology Section, 227 (1992) 455–458
© 1992 Elsevier Science Publishers B.V. All rights reserved 0922-4106/92/$05.00

Author index

Aasen, A., see Waelbroeck, M. (227) 33
Abbracchio, M.P., Fogliatto, G., Paoletti, A.M., Rovati, G.E. and Cattabeni, F.
Prolonged in vitro exposure of rat brain slices to adenosine analogues: selective desensitization of adenosine A_1 but not A_2 receptors (227) 317
Abe, T., see Tokumaru, H. (227) 363
Abrahams, T.P. and Schofield, G.G.
Norepinephrine-induced Ca^{2+} current inhibition in adult rat sympathetic neurons does not require protein kinase C activation (227) 189
Aikawa, J., see Namiki, A. (227) 339
Akil, H., see Mansour, A. (227) 205
Akiyoshi, J., see Lesch, K.P. (227) 233
Aly, M.H.M., see Mohamed, A.S. (227) 181
Anzai, K., see Tokumaru, H. (227) 363
Appenheimer, M., see Ko, Y. (227) 215
Arakawa, M., see Suzuki, S. (227) 395
Aulakh, C.S., see Lesch, K.P. (227) 109
Aulakh, C.S., see Lesch, K.P. (227) 233

Baganoff, M.P., see Davies, M.S. (227) 51
Ball, S.G., see Wintersgill, H.P. (227) 283
Ballesta, J.J. and Orts, A.
Interaction of opioid drugs with human platelet α_2-adrenoceptors (227) 349
Balmforth, A.J., see Wintersgill, H.P. (227) 283
Baraban, J.M., see Bhat, R.V. (227) 447
Basile, A.S., Paul, I.A. and De Costa, B.
Differential effects of cytochrome P-450 induction on ligand binding to σ receptors (227) 95
Bast, A., see Voss, H.-P. (227) 403
Bazan, E., Campbell, A.K. and Rapoport, R.M.
Protein kinase C activity in blood vessels from normotensive and spontaneously hypertensive rats (227) 343
Bergmann, J.S., see Johnson, K.M. (227) 411
Bertacco, S., see Muccioli, G. (227) 293
Bettini, E., see Spreafico, E. (227) 353
Bhat, R.V. and Baraban, J.M.
High basal expression of *zif268* in cortex is dependent on intact noradrenergic system (227) 447
Biagi, M., see Brown, M.C.S. (227) 163
Bidlack, J.M., Saripalli, L.D. and Lawrence, D.M.P.
κ-Opioid binding sites on a murine lymphoma cell line (227) 257
Bitran, J.A., see Manji, H.K. (227) 275
Brands, J., see Stam, N.J. (227) 153
Briggs, R.S.J., see Porter, R.H.P. (227) 83
Brørs, O., see Stokke, M. (227) 379
Brown, M.C.S., Biagi, M., Renzetti, A.R., Rovero, P., Criscuoli, M. and Mizrahi, J.
[^{125}I]His-neurokinin A binds selectively to NK_2 receptors of the B-type in rat small intestine smooth muscle membranes (227) 163
Bruinvels, A.T., Landwehrmeyer, B., Moskowitz, M.A. and Hoyer, D.
Evidence for the presence of 5-HT_{1B} receptor messenger RNA in neurons of the rat trigeminal ganglia (227) 357
Bryson, S.E., see Wintersgill, H.P. (227) 283
Burbach, J.P.H. and Meijer, O.C.
The structure of neuropeptide receptors (227) 1

Cadoni, C. and Gee, K.W.
Complex interactions between the steroid derivative RU 5135 and the $GABA_A$-receptor complex (227) 147
Campbell, A.K., see Bazan, E. (227) 343
Camus, J., see Waelbroeck, M. (227) 33
Caspar Rüegg, J., see Strauss, J.D. (227) 437
Casteels, R., see Missiaen, L. (227) 391
Cattabeni, F., see Abbracchio, M.P. (227) 317
Chan-Yeung, M., see Howard, S. (227) 123
Chang, C.-H., Yu, Z.-N. and Song, D.-L.
Formycin triphosphate as a probe for the ATP binding site involved in the activation of guanylate cyclase (227) 229
Chen, G., see Manji, H.K. (227) 275
Chevillard, C., see Diochot, S. (227) 427
Chiasson, B.J., Hooper, M.L., Murphy, P.R. and Robertson, H.A.
Antisense oligonucleotide eliminates in vivo expression of c-*fos* in mammalian brain (227) 451
Chiavaroli, C., Vacher, P. and Schlegel, W.
Modulation of Ca^{2+} influx by protein phosphorylation in single intact clonal pituitary cells (227) 173
Chiu, A.T., see Leung, K.H. (227) 63
Christophe, J., see Delporte, C. (227) 247
Christophe, J., see Waelbroeck, M. (227) 33
Chuang, D.-M., see Lesch, K.P. (227) 233
Civelli, O., see Mansour, A. (227) 205
Civelli, O., see Seeman, P. (227) 139
Couture, R., see Hasséssian, H. (227) 103
Criscuoli, M., see Brown, M.C.S. (227) 163

Daly, J.W., see Murata, M. (227) 43
Davies, M.S., Baganoff, M.P., Grishin, E.V., Lanthorn, T.H., Volkova, T.M., Watson, G.B. and Wiegand, R.C.
Polyamine spider toxins are potent un-competitive antagonists of rat cortex excitatory amino acid receptors (227) 51
De Costa, B., see Basile, A.S. (227) 95
De Costa, B., see Reith, M.E.A. (227) 417
DeHaven-Hudkins, D.L., Fleissner, L.C. and Ford-Rice, F.Y.
Characterization of the binding of [^{3}H](+)-pentazocine to σ recognition sites in guinea pig brain (227) 371
Delporte, C., Poloczek, P., Tastenoy, M., Winand, J. and Christophe, J.
Atrial natriuretic peptide binds to ANP-R_1 receptors in neuroblastoma cells or is degraded extracellularly at the Ser-Phe bond (227) 247
Demchyshyn, L., see Miller, K.J. (227) 99
De Smedt, H., see Missiaen, L. (227) 391
Diamond, J., see Langlands, J.M. (227) 131
Di Carlo, R., see Muccioli, G. (227) 293
Dijkema, R., see Stam, N.J. (227) 153
Diochot, S., Vittet, D., Mathieu, M.-N., Launay, J.-M. and Chevillard, C.

Effects of endothelins on the human megakaryoblastic cell line MEG-01 (227) 427
Donnell, D., see Voss, H.-P. (227) 403
Droogmans, G., see Missiaen, L. (227) 391
Düsing, R., see Ko, Y. (227) 215

Eiger, S., see Lameh, J. (227) 19
El-Fakahany, E.E., see Mohamed, A.S. (227) 181

Faunce, D.M., see Sawutz, D.G. (227) 309
Fleissner, L.C., see DeHaven-Hudkins, D.L. (227) 371
Fogliatto, G., see Abbracchio, M.P. (227) 317
Folger, W.H., Halushka, P.V., Wilcox, C.S. and Guzman, N.J.
Characterization of rat glomerular thromboxane A_2 receptors: comparison to rat platelets (227) 71
Ford-Rice, F.Y., see DeHaven-Hudkins, D.L. (227) 371
Forray, C., see Mohamed, A.S. (227) 181
Fredholm, B.B., see Kontny, E. (227) 333
Fukazawa, M., see Namiki, A. (227) 339

García-Sáinz, J.A., Macías-Silva, M., Olivares-Reyes, A. and Romero-Avila, M.T.
Histamine activates phosphorylase and inositol phosphate production in guinea pig hepatocytes (227) 325
García-Sáinz, J.A., Romero-Avila, M.T., Olivares-Reyes, J.A. and Macías-Silva, M.
Guinea pig hepatocyte α_{1A}-adrenoceptors: characterization, signal transduction and regulation (227) 239
Gee, K.W., see Cadoni, C. (227) 147
Görg, A., see Ko, Y. (227) 215
Grishin, E.V., see Davies, M.S. (227) 51
Guan, H.-c., see Seeman, P. (227) 139
Gusovsky, F., see Manji, H.K. (227) 275
Gusovsky, F., see Murata, M. (227) 43
Guzman, N.J., see Folger, W.H. (227) 71

Hagelin, E.M., see Stokke, M. (227) 379
Haldeman, B., see Thomsen, C. (227) 361
Halushka, P.V., see Folger, W.H. (227) 71
Harada, J., Ueda, H., Iso, Y. and Satoh, M.
Affinity cross-linked δ-opioid receptor in NG108-15 cells is low molecular weight (25 kDa) and coupled to GTP-binding proteins (227) 301
Hasséssian, H., Prat, A. and Couture, R.
Anaesthetic doses of pentobarbital antagonize phosphatidylinositol hydrolysis induced by substance P or carbachol in the spinal cord and cerebral cortex of the rat (227) 103
Haycock, D., see Sawutz, D.G. (227) 309
Healy, D.P., see Ye, M.-Q. (227) 79
Hill, J.L., see Lesch, K.P. (227) 109
Hill, J.L., see Lesch, K.P. (227) 233
Hirata, Y., see Namiki, A. (227) 339
Holopainen, I., Rau, C. and Wojcik, W.J.
Proposed antagonists at $GABA_B$ receptors that inhibit adenylyl cyclase in cerebellar granule cell cultures of rat (227) 225
Hooper, M.L., see Chiasson, B.J. (227) 451
Houck, W.T., see Sawutz, D.G. (227) 309
Hough, C.J., see Lesch, K.P. (227) 233
Howard, S., Chan-Yeung, M., Martin, L., Phaneuf, S. and Salari, H.
Polyphosphoinositide hydrolysis and protein kinase C activation in guinea pig tracheal smooth muscle cells in culture by leukotriene D_4 involve a pertussis toxin sensitive G-protein (227) 123
Hoyer, D., see Bruinvels, A.T. (227) 357
Huggler, E., see Krause, K.-H. (227) 221

Ishihara, M., see Yamamoto, S. (227) 113
Ishii, A., see Miki, I. (227) 199
Ishikawa, M., see Namiki, A. (227) 339
Iso, Y., see Harada, J. (227) 301

Jacobson, A.E., see Reith, M.E.A. (227) 417
Jiang, H., see Yamamoto, S. (227) 113
Johnson, K.M., Bergmann, J.S. and Kozikowski, A.P.
Cocaine and dopamine differentially protect [^{3}H]mazindol binding sites from alkylation by N-ethylmaleimide (227) 411
Joyce, J.N., see Murray, A.M. (227) 443

Kase, H., see Miki, I. (227) 199
Kato, R., see Yamamoto, S. (227) 113
King, A., see Miller, K.J. (227) 99
Kirino, Y., see Tokumaru, H. (227) 363
Ko, Y., Görg, A., Appenheimer, M., Wieczorek, A.J., Düsing, R., Vetter, H. and Sachinidis, A.
Losartan inhibits the angiotensin II-induced stimulation of the phosphoinositide signalling system in vascular smooth muscle cells (227) 215
Kondo, Y., see Tada, S. (227) 25
Kontny, E., Kvanta, A. and Fredholm, B.B.
Activation of protein kinase C and elevation of cAMP interact synergistically to raise c-Fos and AP-1 activity in Jurkat cells (227) 333
Kozikowski, A.P., see Johnson, K.M. (227) 411
Krause, K.-H., Van Delden, C., Huggler, E., Monod, A. and Lew, D.P.
Differential effects on neutrophil activation of staurosporin and its protein kinase C-selective derivative cgp 41231 (227) 221
Kristensen, P., see Thomsen, C. (227) 361
Kvanta, A., see Kontny, E. (227) 333

Lambrecht, G., see Waelbroeck, M. (227) 33
Lameh, J., Eiger, S. and Sadée, W.
Interaction among μ-opioid receptors and α_2-adrenoceptors on SH-SY5Y human neuroblastoma cells (227) 19
Landwehrmeyer, B., see Bruinvels, A.T. (227) 357
Langlands, J.M. and Diamond, J.
Translocation of protein kinase C in bovine tracheal smooth muscle strips: the effect of methacholine and isoprenaline (227) 131
Lanthorn, T.H., see Davies, M.S. (227) 51
Launay, J.-M., see Diochot, S. (227) 427
Lawrence, D.M.P., see Bidlack, J.M. (227) 257
Lesch, K.P., Aulakh, C.S., Wolozin, B.L., Hill, J.L. and Murphy, D.L.
3-(2-Carboxypiperazin-4-yl)propyl-1-phosphonic acid decreases NMDA receptor mRNA (227) 109
Lesch, K.P., Hough, C.J., Aulakh, C.S., Wolozin, B.L., Tolliver, T.J., Hill, J.L., Akiyoshi, J., Chuang, D.-M. and Murphy, D.L.
Fluoxetine modulates G protein α_s, α_q, and α_{12} subunit mRNA expression in rat brain (227) 233
Leung, K.H., Roscoe, W.A., Smith, R.D., Timmermans, P.B.M.W.M. and Chiu, A.T.
Characterization of biochemical responses of angiotensin II (AT_2) binding sites in the rat pheochromocytoma PC12W cells (227) 63
Lew, D.P., see Krause, K.-H. (227) 221
Liu, Y., see Von Euler, G. (227) 449
Lomasney, J.W., see Schwinn, D.A. (227) 433

Machii, K., see Namiki, A. (227) 339
Macías-Silva, M., see García-Sáinz, J.A. (227) 239
Macías-Silva, M., see García-Sáinz, J.A. (227) 325
Maggi, A., see Spreafico, E. (227) 353
Mais, D.E., True, T.A. and Martinelli, M.J.
Characterization by photoaffinity labelling of the human platelet thromboxane A_2/prostaglandin H_2 receptor: evidence for N-linked glycosylation (227) 267

Maksay, G.
Modification of benzodiazepine receptors supports the distinctive role of histidine residues (227) 57
Manji, H.K., Chen, G., Bitran, J.A., Gusovsky, F. and Potter, W.Z.
Idazoxan down-regulates β-adrenoceptors on C6 glioma cells in vitro (227) 275
Mansour, A., Meng, F., Meador-Woodruff, J.H., Taylor, L.P., Civelli, O. and Akil, H.
Site-directed mutagenesis of the human dopamine D_2 receptor (227) 205
Martin, L., see Howard, S. (227) 123
Martinelli, M.J., see Mais, D.E. (227) 267
Mathieu, M.-N., see Diochot, S. (227) 427
Meador-Woodruff, J.H., see Mansour, A. (227) 205
Meijer, O.C., see Burbach, J.P.H. (227) 1
Meng, F., see Mansour, A. (227) 205
Miki, I., Kase, H. and Ishii, A.
Differences in activities of thromboxane A_2 receptor antagonists in smooth muscle cells (227) 199
Miller, K.J., King, A., Demchyshyn, L., Niznik, H. and Teitler, M.
Agonist activity of sumatriptan and metergoline at the human 5-$HT_{1D\beta}$ receptor: further evidence for a role of the 5-HT_{1D} receptor in the action of sumatriptan (227) 99
Missiaen, L., De Smedt, H., Droogmans, G. and Casteels, R.
2,5-Di-(tert-butyl)-1,4-benzohydroquinone and cyclopiazonic acid decrease the Ca^{2+} permeability of endoplasmic reticulum (227) 391
Mitsui, Y., see Tada, S. (227) 25
Mizrahi, J., see Brown, M.C.S. (227) 163
Mohamed, A.S., Forray, C., Aly, M.H.M. and El-Fakahany, E.E.
Lack of intrinsic activity and significant subtype selectivity of SR 95639A at muscarinic receptors (227) 181
Monod, A., see Krause, K.-H. (227) 221
Moroi, M., see Namiki, A. (227) 339
Moskowitz, M.A., see Bruinvels, A.T. (227) 357
Muccioli, G., Scordamaglia, A., Bertacco, S. and Di Carlo, R.
Effect of S-adenosyl-L-methionine on brain muscarinic receptors of aged rats (227) 293
Mulvihill, E., see Thomsen, C. (227) 361
Murata, M., Gusovsky, F., Yasumoto, T. and Daly, J.W.
Selective stimulation of Ca^{2+} flux in cells by maitotoxin (227) 43
Murphy, D.L., see Lesch, K.P. (227) 109
Murphy, D.L., see Lesch, K.P. (227) 233
Murphy, P.R., see Chiasson, B.J. (227) 451
Murray, A.M., Ryoo, H. and Joyce, J.N.
Visualization of dopamine D_3-like receptors in human brain with [^{125}I]epidepride (227) 443
Mutschler, E., see Waelbroeck, M. (227) 33

Namiki, A., Hirata, Y., Fukazawa, M., Ishikawa, M., Moroi, M., Aikawa, J., Yamaguchi, T. and Machii, K.
Granulocyte-colony stimulating factor stimulates immunoreactive endothelin-1 release from cultured bovine endothelial cells (227) 339
Nishikawa, K., see Yamamoto, S. (227) 113
Niznik, H., see Miller, K.J. (227) 99
Niznik, H.B., see Seeman, P. (227) 139

Okajima, F., see Tada, S. (227) 25
Olijve, W., see Stam, N.J. (227) 153
Olivares-Reyes, A., see García-Sáinz, J.A. (227) 325
Olivares-Reyes, J.A., see García-Sáinz, J.A. (227) 239
Orts, A., see Ballesta, J.J. (227) 349

Paoletti, A.M., see Abbracchio, M.P. (227) 317
Paul, I.A., see Basile, A.S. (227) 95
Phaneuf, S., see Howard, S. (227) 123
Pollio, G., see Spreafico, E. (227) 353
Poloczek, P., see Delporte, C. (227) 247
Porter, R.H.P., Briggs, R.S.J. and Roberts, P.J.
Modulation of [^{3}H]3-((±)-2-carboxypiperazin-4-yl)propyl-1-phosphonic acid ([^{3}H]CPP) binding by ligands acting at the glycine and the polyamine sites of the rat brain NMDA receptor complex (227) 83
Potter, W.Z., see Manji, H.K. (227) 275
Poulsson, C., see Stokke, M. (227) 379
Prat, A., see Hasséssian, H. (227) 103

Rapoport, R.M., see Bazan, E. (227) 343
Rau, C., see Holopainen, I. (227) 225
Reiser, G.
Nitric oxide formation caused by Ca^{2+} release from internal stores in neuronal cell line is enhanced by cyclic AMP (227) 89
Reith, M.E.A., De Costa, B., Rice, K.C. and Jacobson, A.E.
Evidence for mutually exclusive binding of cocaine, BTCP, GBR 12935, and dopamine to the dopamine transporter (227) 417
Rengel, I., Sinowatz, F. and Schulz, R.
Opioid-controlled adenylate cyclase in the guinea-pig myenteric plexus is confined to nerve somata (227) 385
Renzetti, A.R., see Brown, M.C.S. (227) 163
Rice, K.C., see Reith, M.E.A. (227) 417
Roberts, P.J., see Porter, R.H.P. (227) 83
Robertson, H.A., see Chiasson, B.J. (227) 451
Romero-Avila, M.T., see García-Sáinz, J.A. (227) 239
Romero-Avila, M.T., see García-Sáinz, J.A. (227) 325
Roscoe, W.A., see Leung, K.H. (227) 63
Rovati, G.E., see Abbracchio, M.P. (227) 317
Rovero, P., see Brown, M.C.S. (227) 163
Ryoo, H., see Murray, A.M. (227) 443

Sachinidis, A., see Ko, Y. (227) 215
Sadée, W., see Lameh, J. (227) 19
Sakatume, M., see Suzuki, S. (227) 395
Salari, H., see Howard, S. (227) 123
Saripalli, L.D., see Bidlack, J.M. (227) 257
Sato, H., see Suzuki, S. (227) 395
Satoh, M., see Harada, J. (227) 301
Sawutz, D.G., Faunce, D.M., Houck, W.T. and Haycock, D.
Characterization of bradykinin B_2 receptors on human IMR-90 lung fibroblasts: stimulation of $^{45}Ca^{2+}$ efflux by D-Phe^7 substituted bradykinin analogues (227) 309
Schjelderup, L., see Waelbroeck, M. (227) 33
Schlegel, W., see Chiavaroli, C. (227) 173
Schofield, G.G., see Abrahams, T.P. (227) 189
Schulz, R., see Rengel, I. (227) 385
Schwinn, D.A. and Lomasney, J.W.
Pharmacologic characterization of cloned α_1-adrenoceptor subtypes: selective antagonists suggest the existence of a fourth subtype (227) 433
Scordamaglia, A., see Muccioli, G. (227) 293
Seeman, P., Guan, H.-c., Civelli, O., Van Tol, H.H.M., Sunahara, R.K. and Niznik, H.B.
The cloned dopamine D_2 receptor reveals different densities for dopamine receptor antagonist ligands. Implications for human brain positron emission tomography (227) 139
Sinowatz, F., see Rengel, I. (227) 385
Smith, R.D., see Leung, K.H. (227) 63
Song, D.-L., see Chang, C.-H. (227) 229
Spreafico, E., Bettini, E., Pollio, G. and Maggi, A.
Nucleotide sequence of estrogen receptor cDNA from Sprague-Dawley rat (227) 353
Stam, N.J., Van Huizen, F., Van Alebeek, C., Brands, J., Dijkema, R., Tonnaer, J.A.D.M. and Olijve, W.
Genomic organization, coding sequence and functional expression of human 5-HT_2 and 5-HT_{1A} receptor genes (227) 153

Stokke, M., Hagelin, E.M., Poulsson, C. and Brørs, O.
Ca^{2+} modulates the inhibition of (+)-[^{3}H]isradipine binding by amiloride and quinacrine (227) 379
Strauss, J.D., Zeugner, C. and Caspar Rüegg, J.
The positive inotropic calcium sensitizer EMD 53998 antagonizes phosphate action on cross-bridges in cardiac skinned fibers (227) 437
Strohmann, C., see Waelbroeck, M. (227) 33
Sugai, K., see Suzuki, S. (227) 395
Sunahara, R.K., see Seeman, P. (227) 139
Suzdak, P.D., see Thomsen, C. (227) 361
Suzuki, S., Sugai, K., Sato, H., Sakatume, M. and Arakawa, M.
Inhibition of active oxygen generation by dipyridamole in human polymorphonuclear leukocytes (227) 395

Tacke, R., see Waelbroeck, M. (227) 33
Tada, S., Okajima, F., Mitsui, Y., Kondo, Y. and Ui, M.
P_2 purinoceptor-mediated cyclic AMP accumulation in bovine vascular smooth muscle cells (227) 25
Tastenoy, M., see Delporte, C. (227) 247
Tastenoy, M., see Waelbroeck, M. (227) 33
Taylor, L.P., see Mansour, A. (227) 205
Teitler, M., see Miller, K.J. (227) 99
Thomsen, C., Kristensen, P., Mulvihill, E., Haldeman, B. and Suzdak, P.D.
L-2-Amino-4-phosphonobutyrate (L-AP4) is an agonist at the type IV metabotropic glutamate receptor which is negatively coupled to adenylate cyclase (227) 361
Timmermans, P.B.M.W.M., see Leung, K.H. (227) 63
Tokumaru, H., Anzai, K., Abe, T. and Kirino, Y.
Purification of the cardiac 1,4-dihydropyridine receptor using immunoaffinity chromatography with a monoclonal antibody against the $\alpha_2\delta$ subunit of the skeletal muscle dihydropyridine receptor (227) 363
Tolliver, T.J., see Lesch, K.P. (227) 233
Tonnaer, J.A.D.M., see Stam, N.J. (227) 153
True, T.A., see Mais, D.E. (227) 267

Ueda, H., see Harada, J. (227) 301
Ui, M., see Tada, S. (227) 25

Vacher, P., see Chiavaroli, C. (227) 173
Van Alebeek, C., see Stam, N.J. (227) 153
Van Delden, C., see Krause, K.-H. (227) 221
Van Huizen, F., see Stam, N.J. (227) 153
Van Tol, H.H.M., see Seeman, P. (227) 139
Vetter, H., see Ko, Y. (227) 215
Vittet, D., see Diochot, S. (227) 427
Volkova, T.M., see Davies, M.S. (227) 51
Von Euler, G. and Liu, Y.
The glutamate- and glycine-induced reduction of [^{3}H]MK-801 binding in the presence of Mg^{2+} is reversed at low pH (227) 449
Voss, H.-P., Donnell, D. and Bast, A.
Atypical molecular pharmacology of a new long-acting β_2-adrenoceptor agonist, TA 2005 (227) 403

Waelbroeck, M., Camus, J., Tastenoy, M., Mutschler, E., Strohmann, C., Tacke, R., Schjelderup, L., Aasen, A., Lambrecht, G. and Christophe, J.
Stereoselective interaction of procyclidine, hexahydro-difenidol, hexbutinol and oxyphencyclimine, and of related antagonists, with four muscarinic receptors (227) 33
Wang, J.C., see Yamamoto, S. (227) 113
Warburton, P., see Wintersgill, H.P. (227) 283
Watson, G.B., see Davies, M.S. (227) 51
Wieczorek, A.J., see Ko, Y. (227) 215
Wiegand, R.C., see Davies, M.S. (227) 51
Wilcox, C.S., see Folger, W.H. (227) 71
Winand, J., see Delporte, C. (227) 247
Wintersgill, H.P., Warburton, P., Bryson, S.E., Ball, S.G. and Balmforth, A.J.
Characterization of the angiotensin II receptor expressed by the human hepatoma cell line, PLC-PRF-5 (227) 283
Wojcik, W.J., see Holopainen, I. (227) 225
Wolozin, B.L., see Lesch, K.P. (227) 109
Wolozin, B.L., see Lesch, K.P. (227) 233

Yamaguchi, T., see Namiki, A. (227) 339
Yamamoto, S., Jiang, H., Nishikawa, K., Ishihara, M., Wang, J.C. and Kato, R.
Protein kinase C-dependent and -independent actions of a potent protein kinase C inhibitor, staurosporine (227) 113
Yasumoto, T., see Murata, M. (227) 43
Ye, M.-Q., Yuan, L.-X. and Healy, D.P.
Effects of intracerebroventricular captopril on angiotensin-converting enzyme and angiotensinogen mRNA levels in rat brain (227) 79
Yu, Z.-N., see Chang, C.-H. (227) 229
Yuan, L.-X., see Ye, M.-Q. (227) 79

Zeugner, C., see Strauss, J.D. (227) 437